

## CLAYTON RALLY MAPP 13415

contains 50 g/l tepraloxym in an emulsifiable concentrate with solvent naphtha

**For the control of certain grass weeds in the listed broad-leaved crops and on green cover on land not being used for crop production.**



**HARMFUL**

**Harmful: may cause lung damage if swallowed**  
**Irritating to skin**  
**Possible risk of harm to the unborn child**  
**Possible risk of impaired fertility**  
**Limited evidence of a carcinogenic effect**  
**Repeated exposure may cause skin dryness or cracking**



**DANGEROUS FOR THE ENVIRONMENT**

**Toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment**

The Control of Substances Hazardous to Health (COSHH) Regulations may apply to the use of this product at work.

### IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL HERBICIDE			
Crop	Maximum individual dose of product	Maximum number of applications	Latest time of application
Winter oilseed rape	1 l/ha	1 per crop	Before end-November or before the crop has 9 true leaves, whichever first occurs.
Linseed	1.5 l/ha	1 per crop	Before the flower buds are visible.
Sugar beet Fodder beet Field bean	1.5 l/ha	1 per crop	8 weeks before harvest.
Combining pea Vining pea	1.5 l/ha	1 per crop	5 weeks before harvest.
Leeks	1.5 l/ha	1 per crop	4 weeks before harvest.
Bulb onions	Or 0.75 l/ha	2 per crop	
Carrots	1.5 l/ha	1 per crop	3 weeks before harvest
Cabbage	1.5 l/ha	1 per crop	Before head formation (when the two youngest leaves do not unfold).
Cauliflower	1.5 l/ha	1 per crop	Before head formation.
Green cover on land not being used for crop production	1.5 l/ha	1 per year	See 'Other specific restrictions'.
<b>Other specific restrictions</b>			
<ol style="list-style-type: none"> <li>1. For linseed, sugar beet, fodder beet and green cover on land not being used for crop production, applications are prohibited between 1 November and 31 March.</li> <li>2. For field bean, combining pea, vining pea, leek, bulb onion, carrot, cabbage and cauliflower, applications are prohibited between 1 November and 1 March.</li> <li>3. Where this product is used on green cover on land not being used for crop production, a full green cover must be established before the pesticide is applied.</li> <li>4. Where this product is used on green cover on land not being used for crop production, treated plants must not be grazed by livestock or harvested for human or animal consumption.</li> <li>5. To avoid the build-up of resistance do not apply products containing an ACCase inhibitor herbicide more than twice to winter oilseed rape, linseed, sugar beet, fodder beet, field beans, combining peas, vining peas, carrots, cabbage, cauliflower and green cover on land not being used for crop production. In addition, in these crops do not use this product in mixture or sequence with any other product containing tepraloxym.</li> </ol>			
<b>READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.</b>			

## **SAFETY PRECAUTIONS**

### **Operator protection**

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment:

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate and when handling contaminated surfaces.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Do not breathe spray.

WASH CONCENTRATE from skin or eyes immediately.

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before eating and drinking and after work.

This product must be applied only via conventional ground sprayers; do not apply via a knapsack sprayer.

IF SWALLOWED, do not induce vomiting. Seek medical advice and show this container or label.

In case of accident or if you feel unwell seek medical advice immediately (show the label where possible).

### **Environmental protection**

Do not contaminate water with the product or its container. Do not clean application equipment near surface water. Avoid contamination via drains from farmyards and roads.

### **Storage and disposal**

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

This material and its container must be disposed of in a safe way.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of the container safely.

### **To avoid risks to man and the environment, comply with the instructions for use.**

Safety data sheet available for professional user on request.

**Clayton Plant Protection (UK) Ltd.,  
Unit F10, Bracetown Business Park  
CLONEE, Co. Meath  
Ireland.**

Tel: (00 353) 1 8210127

Contents: **1 litre**

STORE IN A COOL DRY PLACE

PROTECT FROM FROST

SHAKE BEFORE POURING

Batch No:

This product is approved under The Plant Protection Products Regulations (as amended).

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (contains solvent naphtha). UN 3082; Class 9; Packing Group III.
--

**Conditions of Supply:** all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

## DIRECTIONS FOR USE

**IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.**

### MODE OF ACTION

Clayton Rally is a leaf-absorbed graminicide. Susceptible grass weeds need to have at least 2-4 fully expanded leaves at application – see RECOMMENDATIONS table for precise conditions. Weeds not emerged at application are not controlled. Growth of susceptible grass weeds is stopped within a few days of application; they are controlled most rapidly when the weather is warm and the soil moist. During cool weather, weed control is slowed, most particularly when applied in late autumn to winter oilseed rape. Under favourable conditions break down at the growing point is apparent after 2-3 weeks. Vigorous crop competition is an essential requirement for optimum weed control, especially when the grass weeds being treated are at the later growth stages of their treatable range. Some recovery might occur in thin crops offering little competition or if lack of soil moisture is limiting growth.

### RECOMMENDATIONS - WEEDS

Spray before grass weeds or sown covers, e.g. barley protecting sugar beet, become unduly competitive with the crop.

For common couch, cultivations should not be conducted in the period before treatment but may be resumed 14-21 days after application depending upon the intervening growing conditions. Delay any cultivation for at least 21 days when growing conditions have been poorer. Note however that cultivation may induce fresh weed germination.

The long-term effectiveness of this product on common couch and creeping bent is not known. Maximum efficacy is gained when the underground stems to plough depth have been chopped or cut by cultivations prior to planting.

Aim to treat weeds when they are most susceptible, that is when small before tillering; a lower level of weed control may be achieved when the time of application is sub-optimal.

Target grass	Dose	Spraying period and remarks
Black-grass – in winter oilseed rape and field beans only	1 l/ha (14 fl.oz/ac)	Autumn application: from 2 fully-expanded leaves up to 5 tillers. Spring application: from 2 fully-expanded leaves to the end of tillering.
Sown barley cover Volunteer barley Wild-oat	1 l/ha (14 fl.oz/ac)	From 2 fully-expanded leaves to the end of tillering.
Volunteer wheat Sterile Brome	1 l/ha (14 fl.oz/ac)	From 2 fully-expanded leaves up to 5 tillers.
Annual meadow-grass	1.5 l/ha (21 fl.oz/ac)	Autumn application: from 2 fully-expanded leaves up to 2 tillers. Spring application: from 2 fully-expanded leaves up to 4 tillers.
Annual meadow-grass - split dose recommendation for bulb onions and leeks only.	Two applications each of 0.75 l/ha (10½ fl.oz/ac)	Provided that black-grass, Italian rye-grass or wild-oats are not also present (in which case revert to the full dose application above), the total dose may be applied in two applications within the spraying period given for the full dose above.
Couch, common	1.5 l/ha (21 fl.oz/ac)	When all the shoots have emerged and are about 15 cm long. Use specifically against couch only as an emergency in-crop treatment as kill of underground stems cannot be assured.

### WEED RESISTANCE

Strains of some annual grasses, e.g. black-grass, wild-oat and Italian rye-grass, have developed resistance to herbicides which may lead to poor control. A strategy for preventing and managing such resistance should be adopted. Guidelines have been produced by the Weed Resistance Action Group and copies are available from the HGCA, CPA, your distributor, crop advisor or product manufacturer.

This product contains tepraloxymid which is an ACCase inhibitor; also classified by the Herbicide Resistance Action Committee as 'Group A'.

Use only as part of a resistance management strategy that includes cultural methods of control and does not use ACCase inhibitors as the sole chemical method of grass-weed control.

Applying a second product containing an ACCase inhibitor to a crop will increase the risk of resistance development; only use a second ACCase inhibitor to control different weeds at a different timing.

If the presence of herbicide resistant strains is known or suspected do not apply less than the full recommended dose of Clayton Rally for the situation.

- Always follow WRAG guidelines for preventing and managing herbicide resistant grass-weeds.
- Do not use Clayton Rally or any other ACCase inhibitor as the sole means of grass weed control in successive crops.
- Use grass-weed herbicides with different modes of action throughout the cropping rotation.
- To reduce the risk of developing resistance, applications should be made to young, actively growing weeds.
- Use mixtures of herbicides or sequences of herbicides with different modes of action within individual crops or successive crops.
- Use crop rotation and other cultural control measures to prevent and manage herbicide resistant grass weeds.
- Monitor treatment efficacy and seek professional advice if areas of poor weed control occur that appear otherwise inexplicable.

#### RECOMMENDATIONS – CROPS

Crop	Time of application and remarks
Winter oilseed rape	Spray from one expanded true-leaf until closure of the crop canopy or before end-November or before the plants have 9 true-leaves, whichever first occurs. Do not treat crops lacking vigour.
Sugar beet Fodder beet	Spray from 1 April from two expanded true-leaves to before the crop leaves cover the weeds. Do not treat later than 8 weeks before harvest.
Winter field bean	Spray in the spring from 1 March from 3 leaves until 6 internodes can be counted.
Spring field bean	Spray from 1 March from two leaves until the first side shoots appear.
Cabbage	Spray from 1 March from four expanded true-leaves to before the start of head formation (when the two youngest leaves do not unfold).
Cauliflower	Spray from 1 March from two expanded true-leaves to before the start of head formation.
Combining peas Vining peas	Spray from 1 March from two nodes to before the crop leaves cover the weeds and at least 5 weeks before harvest. Treat winter-sown peas only in the spring. A satisfactory crystal violet test for leaf wax must be conducted before treatment.
Bulb onions Leeks	Spray from 1 March from three visible true-leaves to before the crop leaves cover the weeds and at least 4 weeks before harvest. Do not treat crops deficient in leaf wax. Two split-dose treatments may be used for the control of annual meadow-grass in these crops; see RECOMMENDATIONS-WEEDS.
Carrots	Spray from 1 March from one fully expanded true-leaf to before the crop leaves cover the weeds and at least 3 weeks before harvest.
Linseed	Spray from 1 April from two expanded leaves to before the flower buds become visible.
Green cover on land not being used for crop production (set-aside)	Spray between 1 April and 31 October from the green cover being fully established. The initial green cover must be composed of sufficient tolerant species so that an adequate green cover remains after removal of the weed grasses. Check for compliance with any grant-aided scheme rules before treating. Livestock must not graze treated areas nor must any area be harvested for human or animal consumption.

Ensure that application is made before closure of the crop canopy so that weeds are not shielded from the spray by the crop foliage.

The leaf wax is satisfactory if less than 5 % of the upper leaf surface and less than 10% of the lower leaf surface of typical leaves retain solution when dipped into 1% crystal violet solution.

#### COMPATIBILITY IN A PROGRAMME

Clayton Rally can normally be accommodated within most weed control programmes for the recommended crops provided that an interval of at least 14 days either side is allowed between treatments. The leaf wax of crops must be intact before treatment; if in doubt delay an application.

### **CROP, SOIL AND WEATHER FACTORS**

Best results are obtained when the weather is warm and the soil is moist, encouraging active growth. Do not apply to crops under any growing stress such as drought, waterlogging, physical damage, pesticide toxicity, improper nutrition or other disorder until the stresses have been relieved and normal growth has resumed.

Do not apply during periods of cold weather, frost or snow or if cold weather has been forecast.

Do not apply to wet foliage or if rain is forecast within 4 hours.

Do not apply to crops under stress during hot weather.

### **SPRAY MIXING**

Shake the product in its container thoroughly before pouring. Half-fill the spray tank with clean water and put under agitation. Mix in the required volume of Clayton Rally through the top filter or filling device. Top up the tank with water and keep under agitation until sprayed out. Spray immediately after mixing. This product *does not require* the addition of adjuvant oil.

### **APPLICATION**

Apply Clayton Rally through a mounted or drawn conventional hydraulic sprayer. Apply the recommended dose as a FINE or MEDIUM (BCPC) spray in 100-200 l/ha (9-18 gal/ac) water, but employ at least 200 l/ha (18 gal/ac) water when treating perennial grasses. Use the highest spray volume in the most dense crops or when weeds are numerous or large. Avoid spray drift onto nearby crops, especially onto cereals and grasses that are particularly susceptible to damage. Do not overlap spray swaths.

### **PROCESSED CROPS**

Consult processors before treating crops being grown for processing.

### **FOLLOWING CROPS**

Some restrictions apply to the sowing of certain following crops in the event of crop failure for any reason or after normal harvest. The conditions listed below apply between the last application of Clayton Rally and the sowing or planting of the following crop:

Broad-leaved crops	These crops may follow as and how required.
Wheat and barley	Minimum 2 weeks interval and after normal seedbed cultivations.
Forage maize and Italian rye-grass	Minimum 8 weeks interval with ploughing to 20cm and soil mixing cultivations.
Cereals and grasses not named above	These crops must not be planted as the immediate following crop.

### **AFTER USE**

Immediately after use, thoroughly clean the sprayer with water and a cleaning agent recommended for the cleaning of spraying machines. Susceptible crops could be damaged by traces of tepraloxydim left in the sprayer.

### **MAXIMUM RESIDUE LEVELS**

The following maximum residue levels must not be exceeded:

<b>Crop</b>	<b>Maximum residue level (mg/kg)</b>
Oilseed rape	1.0
Linseed/Flax	1.0
Field bean (dry)	0.5
Combining pea (dry)	0.5
Vining pea (fresh without pod)	0.2
Leek	0.2
Bulb onion	0.5
Carrot	0.5
Cabbage	1.0
Cauliflower	0.5