

CLAYTON SPIGOT

MAPP 11560

Contains 750 g/l (79.5% w/w) fenpropimorph with cyclohexanone
in an emulsifiable concentrate

Fungicide for the control of certain foliar diseases of the listed cereals.



HARMFUL

Irritating to skin
Possible risk of harm to the unborn child



**DANGEROUS FOR THE
ENVIRONMENT**

**Very toxic to aquatic organisms, may cause long-term
adverse effects in the aquatic environment**

The Control of Substances Hazardous to Health (COSHH) Regulations may apply to the use of this product at work.

IMPORTANT INFORMATION

FOR USE ONLY AS AN AGRICULTURAL FUNGICIDE

Crop	Maximum individual dose of product	Maximum number of treatments	Latest time of application	Other specific restrictions
Winter wheat * Winter barley * Winter oats	1 l/ha	3 per crop see Other specific restrictions	5 weeks before harvest	Maximum 2 applications between 1 January and harvest
Rye Triticale	or 0.75 l/ha	4 per crop see Other specific restrictions	5 weeks before harvest	Maximum 3 applications between 1 January and harvest
Spring wheat * Spring barley * Spring oats	1 l/ha	2 per crop	5 weeks before harvest	-

* includes crops undersown with grass or clovers

READ THE LABEL BEFORE USE. USING THIS PRODUCT IN A MANNER THAT IS INCONSISTENT WITH THE LABEL MAY BE AN OFFENCE. FOLLOW THE CODE OF PRACTICE FOR USING PLANT PROTECTION PRODUCTS.

SAFETY PRECAUTIONS

Operator protection

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.

WEAR SUITABLE PROTECTIVE GLOVES when handling the concentrate.

However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

Wear suitable protective clothing and gloves.

AVOID ALL CONTACT WITH SKIN OR EYES.

WASH CONCENTRATE from skin or eyes immediately.

DO NOT BREATHE SPRAY.

WHEN USING, DO NOT EAT, DRINK OR SMOKE.

WASH HANDS AND EXPOSED SKIN before meals and after work.

If swallowed, seek medical advice immediately and show this container or label.

Environmental protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

This material and its container must be disposed of in a safe way.

RINSE CONTAINER THOROUGHLY by using an integrated pressure rinsing device or manually rinsing three times. Add washings to sprayer at time of filling and dispose of the container safely.

To avoid risks to man and the environment, comply with the instructions for use.

Safety data sheet available for professional user on request.

This product is approved under the Control of Pesticides Regulations 1986.

Clayton Plant Protection (UK) Ltd.
Unit F10, Bracetown Business Park
CLONEE, Co. Meath
Ireland.

Tel: (00 353) 1 8210127

Contents: **5 litres**

PROTECT FROM FROST

Keep in a dry, cool pesticide store

Batch No:

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S. (contains fenpropimorph). UN 3082, Class 9, Packing group III
--

Conditions of Supply: all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Clayton Spigot is a fungicide of the morpholine group and has contact and systemic activity. Apply immediately disease becomes active, particularly when yellow and brown rusts are amongst the target diseases or the weather favours disease spread.

WINTER AND SPRING WHEAT

Disease	Timing
Powdery mildew	Spray as soon as active mildew is visible, at least before infection reaches 2% on the lower leaves. Repeat if necessary.
Brown rust	Immediately brown rust appears on the upper leaves. If a severe attack threatens, spray at flag leaf emerged and again at ear emerged but before flowering.
Yellow rust	Treat on sight of the disease, at least before infection reaches 1% on any leaf. If necessary repeat 2-3 weeks later. Timely treatment to prevent the disease becoming established gives the best results.

TRITICALE

Disease	Timing
Brown rust	Immediately brown rust appears on the upper leaves. If a severe attack threatens, spray at flag leaf emerged and again at ear emerged but before flowering.
Yellow rust	Treat on sight of the disease, at least before infection reaches 1% on any leaf. If necessary repeat 2-3 weeks later. Timely treatment to prevent the diseases becoming established gives the best results.

WINTER BARLEY

Disease	Timing
Powdery mildew	Autumn: spray as soon as mildew becomes active, particularly on the lighter soils or in thin crops. Spring and summer: spray as soon as active mildew is visible, at least before infection reaches 2% on the lower leaves. Repeat if necessary.
Leaf blotch (<i>Rhynchosporium</i>) for light infections	Spray as soon as possible if leaf blotch is spreading to the uppermost three leaves, particularly around the time of flag leaf emergence.
Brown rust	Immediately brown rust appears on the upper leaves of susceptible varieties or as soon as it begins to spread in less susceptible varieties. If a severe attack threatens, spray at flag leaf emerged and again at ear emerged but before flowering.
Yellow rust	Treat on sight of the disease, at least before infection reaches 1% on any leaf. If necessary repeat 2-3 weeks later. Timely treatment to prevent the disease becoming established gives the best results.

SPRING BARLEY

Disease	Timing
Powdery mildew	Spray as soon as active mildew is visible, at least before infection reaches 2% on the lower leaves. Repeat if necessary.
Leaf blotch (<i>Rhynchosporium secalis</i>) for light infections	Spray immediately leaf blotch is seen in susceptible varieties. In less susceptible varieties, spray as soon as possible if leaf blotch is spreading to the uppermost three leaves, particularly around the time of flag leaf emergence.
Brown rust	Immediately brown rust appears on the upper leaves of susceptible varieties or as soon as it begins to spread in less susceptible varieties. If a severe attack threatens, spray at flag leaf emerged and again at ear emerged but before flowering.
Yellow rust	Treat on sight of the disease, at least before infection reaches 1% on any leaf. If necessary repeat 2-3 weeks later. Timely treatment to prevent the disease becoming established gives the best results.

RYE AND WINTER AND SPRING OATS

Disease	Timing
Powdery mildew	Spray as soon as active mildew is visible, at least before infection reaches 2% on the lower leaves. Repeat if necessary.

RATE OF APPLICATION

The full application rate to control the range of diseases is 1 litre Clayton Spigot per ha per application. However powdery mildew in wheat and barley and brown rust in barley can be controlled satisfactorily by Clayton Spigot at 0.75 l/ha.

APPLICATION

Listed cereals: apply Clayton Spigot as a MEDIUM spray (BCPC definition) at 2-3 bar (30-40 psi) in 220-450 litres of water per ha.

Use the higher water volume when crop growth is dense or the disease pressure is high. Avoid spray drift onto neighbouring crops.

WEATHER AND CROP CONDITIONS

Do not treat crops under growing stresses however caused. Spray onto dry leaves when rain is not imminent. The spray becomes rainfast on cereals about 2 hours after application. Do not apply during frosty weather or when temperatures are high.

COMPATIBILITY

Clayton Spigot is physically compatible for co-application with certain approved pesticides when used in the listed cereals in the spring or summer. Please contact your distributor for further information. When tank-mixing follow the Directions for Use of the partner product together with those of this label.

MIXING

Half-fill the sprayer tank with clean water. With the contents of the spray tank under agitation, add the required quantity of Clayton Spigot. Maintain agitation whilst topping up the tank to the required level and until the completion of spraying. Spray immediately after mixing. When tank-mixing, mix the partner product in the spray tank before mixing in the Clayton Spigot. Wash all equipment thoroughly immediately after use.

UNDERSOWN CROPS

Undersown clovers may be subject to some leaf spotting.
