

CLAYTON VERT

MAPP 12570

contains 250 g/l iprodione in a suspension concentrate

Fungicide for use against certain diseases of managed amenity turf.



HARMFUL

IRRITATING TO EYES AND SKIN
Limited evidence of a carcinogenic effect



DANGEROUS FOR THE ENVIRONMENT

Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment

The Control of Substances Hazardous to Health (COSHH) Regulations may apply to the use of this product at work.

COMPLIANCE WITH THE FOLLOWING CONDITIONS OF USE AND ALL SAFETY PRECAUTIONS MARKED ❖ IS A LEGAL REQUIREMENT

FOR USE ONLY AS A HORTICULTURAL FUNGICIDE

Situation: Managed amenity turf
Maximum individual dose: 1 litre product per 250 m²

READ ALL OTHER SAFETY PRECAUTIONS AND DIRECTIONS FOR USE BEFORE USE

SAFETY PRECAUTIONS

Operator protection

- ❖ Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment.
- ❖ WEAR SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.
- ❖ However, engineering controls may replace personal protective equipment if a COSHH assessment shows they provide an equal or higher standard of protection.

WASH CONCENTRATE from skin or eyes immediately.

WHEN USING DO NOT EAT, DRINK OR SMOKE.

TAKE OFF IMMEDIATELY all contaminated clothing.

WASH HANDS before eating and drinking and after work.

IF YOU FEEL UNWELL, seek medical advice immediately (show the label where possible).

Environmental protection

DO NOT CONTAMINATE SURFACE WATERS OR DITCHES with chemical or used container.

Storage and disposal

KEEP OUT OF REACH OF CHILDREN.

KEEP AWAY FROM FOOD, DRINK AND ANIMAL FEEDING STUFFS.

KEEP IN ORIGINAL CONTAINER, tightly closed, in a safe place.

This material and its container must be disposed of in a safe way.

WASH OUT CONTAINER THOROUGHLY, empty washings into spray tank, and dispose of safely.

To avoid risks to man and the environment, comply with the instructions for use.

Safety data sheet available for professional user on request.

Clayton Plant Protection (UK) Ltd.,
Unit F10
Bracetown Business Park
CLONEE
Co. Meath
Ireland.
Tel: (00 353) 1 8210127

Contents: **5 litres**

PROTECT FROM FROST
SHAKE THOROUGHLY BEFORE USE

Batch No:

This product is approved under the Plant Protection Products Regulations (as amended).

ENVIRONMENTALLY HAZARDOUS SUBSTANCE, LIQUID, N.O.S., (contains iprodione). UN 3082; Class 9; Packing Group III.
--

Conditions of Supply: all goods supplied by us are of high quality and we believe them to be correct but, as we cannot exercise control over their storage, handling, mixing or use, or weather conditions before, during and after application which may affect the performance of the goods, all conditions and warranties, statutory or otherwise, as to the quality or fitness for any purpose of our goods are excluded, and no responsibility will be accepted by us or resellers for any failure in performance, damage or injury whatsoever arising from their storage, handling, application or use. These conditions cannot be varied by our staff or agents whether or not they supervise or assist in the use of such goods.

DIRECTIONS FOR USE

IMPORTANT: This information is approved as part of the Product Label. All instructions within this section must be read carefully in order to obtain safe and successful use of this product.

Clayton Vert is a dicarboximide fungicide with protectant and some curative activity against certain diseases affecting turf.

Disease problem	Timing and remarks	Dose
Dollar spot	Treat when disease is first seen and repeat	20 l/ha in at least 500
Fusarium patch	as necessary. If disease is anticipated treat	l/ha water
Red thread	at monthly intervals.	(200 ml in at least 5
Brown patch	Moderate control of these diseases is	litres water per 100 m ²)
Melting-out	obtained from applications timed as above.	
Grey snow mould	-	

Note: Brown patch, melting-out and grey snow mould are not common in the UK; evidence for efficacy is from non-UK sources. If grey snow mould is the main target disease, the increased dose of 40 l/ha in at least 500 l/ha water is recommended.

FUNGICIDAL EFFICACY

Where strains of disease with reduced sensitivity to the dicarboximide fungicides occur or develop, the level of disease control obtained with Clayton Vert may be reduced and a fungicide with a different mode of action should be used. Alternate use of fungicides with different modes of action may minimise the occurrence of less sensitive strains.

MIXING AND APPLICATION

Shake the container thoroughly before opening. Half-fill the sprayer tank with clean water. With the contents of the spray tank under agitation, add the required quantity of Clayton Vert. Maintain agitation whilst topping up the tank to the required level and until the completion of spraying. Use immediately after mixing. Wash all equipment thoroughly immediately after use.

A knapsack sprayer may be used for small areas. Achieve thorough coverage of the turf. Avoid spray drift.

Treat during a mild period, free of frost. Do not apply to wet grass or when rain is expected. The spray should be dry on the leaves before sunset. Do not treat drought stricken turf or turf under any other kind of stress affecting normal growth. Turf treated whilst under stress, especially during hot or very cold weather, may become yellow for a few days.

AFTER-MANAGEMENT

Avoid mowing for at least 24 hours after application. Preferably do not collect the grass from the first mowing after application.